Eocene of southern British Columbia. 197–209
Diatom, *Neidium rudimentarum* 33–42
Dirca spp. (Thymelaeaceae) reproduce from rhizomes 291–294
Djan-Chékar, Nathalie, Luc Brouillet, Claudia Hanel, Stuart Hay, and John E. Maunder. Vascular plants of the Island of Newfoundland, Canada: Recent additions and rediscoveries. 167–177 (Note)
Draba glabella 55–59; key to New England *Draba* 57–58
Draba glabella (Brassicaceae), new to Maine. 55–59 (New England Note)
Dunwiddie, Peter W. 326–346

Emison, Mary Rose Castelli 273–286
Endangered plants 97–117
Endemics 124–147
Eocene 197–209
Ericaceae 370–376
Eurybia, chromosome numbers 253–272

Farnsworth, Elizabeth J. Patterns of plant invasions at sites with rare plant species throughout New England. 97–117
Flora of Morningside Nature Center, Alachua County, Florida 226–252
Floras of Pequawket and Heath Pond Bogs, Ossipee, New Hampshire. 359–366 (New England Note)
Florida 226–252
Floristic composition and phytodiversity in the wetlands along the base of the Bolivian Andes 1–32
Floristic survey of the forested parts of the United Plant Savers' ethnobotanical sanctuary, Meigs County, Ohio. A. 148–166
Floristics and distribution of vernal pools on the Columbia Plateau of eastern Washington. 326–346
Floristics of inland sandy acid pondshores in the lower Connecticut River Valley 66–76
Floristics: Florida 226–252; southeastern U.S. 305–325
Flowering versus vegetative growth in *Maianthemum canadense* 210–225
Forest community 273–286
Fossil *Neviusia* leaves (Rosaceae: Kerrieae) from the lower-middle Eocene of southern British Columbia. 197–209

Gaylussacia brachycera in North Carolina 370–376; distribution map 371; synonymy 374
Ganger, Michael T. Inferring genet development: Interplay of meristem commitment and genet integrity. 210–225
Genet 210–225
Gentiana quinquefolia var. *occidentalis* (Gentianaceae) 367–369
Gentianaceae 367–369
Genus *Physalis* (Solanaceae) in the southeastern United States. The. 305–325
Gesneriaceae 43–51
Graves, William R. Confirmation that *Dirca* spp. (Thymelaeaceae) reproduce from rhizomes. 291–294 (Note)
Greene, Craig William 77–79 (In Memoriam)

Haines, Arthur 52–54
Haines, Arthur, Andrew Cutko, and Dan Sperduto. Records for *Carex rostrata* (Cyperaceae) in New England. 287–290 (New England Note)
Haines, Arthur. *Draba glabella* (Brassicaceae), new to Maine. 55–59 (New England Note)
Hamilton, Paul B., Peter A. Siver, and Charles W. Reimer. *Neidium rudimentarum* (Bacillariophyceae), a rare species with novel rudimentary canals and areolar openings from the temperate region of North America. 33–42
Hampden Co., Massachusetts 66–76
Hanel, Claudia 167–177
Hay, Stuart 167–177
Hemlock wooly adelgid (*Adelges tsugae*) 273–286

Hemlock-White Pine forest in western Massachusetts 273–286
Hispaniola 124–147
Howard, Richard Alden 178–184 (In Memoriam)

IN MEMORIAM:
Craig William Greene 1949–2003. 77–79
Richard Alden Howard 1917–2003. 178–184
Inferring genet development: Interplay of meristem commitment and genet integrity. 210–225
Inland sandy acid pondshores in the lower Connecticut River Valley, Hampden County, Massachusetts. 66–76 (New England Note)
Invasibility 97–117
Invasive species, New England 97–117
Inventory of the vascular flora of Morningside Nature Center, Alachua County, Florida. An, 226–252

Jahr, Kassandra J. and Garrett E. Crow. Noteworthy plants in created wetlands in southeastern New Hampshire. 60–65 (New England Note)
Jaltomata andersonii, sp. nov., Peru 118–123
Judd, Walter S. 226–252
Judd, Walter S. and Darin S. Penneys. Taxonomic studies in the Miconieae (Melastomataceae). VIII. A revision of the species of the *Miconia desportesii* complex on Hispaniola. 124–147

Kabat, Cathleen A., Steven M. Kabat, and Walter S. Judd. An inventory of the vascular flora of Morningside Nature Center, Alachua County, Florida. 226–252
Kabat, Steven M. 226–252
Kerrieae 197–209
Keys: *Draba* 55–59; *Miconia* 124–147; *Physalis* 305–325; *Pseudosmodingium* 347–358
Kriebel, Ricardo. Two new species and one new name in the Gesneriaceae from Costa Rica. 43–51

Leiva G., Segundo 118–123
Leptotypification of the name *Gentiana quinquefolia* var. *occidentalis* (Gentianaceae). 367–369 (Note)
Liliaceae 210–225

Maianthemum canadense 210–225
Maine: *Carex rostrata* 287–290; *Carex typhina* 52–54; *Draba glabella* 55–59
Martin, Elizabeth and Philip D. Cantino. A floristic survey of the forested parts of the United Plant Savers' ethnobotanical sanctuary, Meigs County, Ohio. 148–166
Massachusetts 66–76, 273–286
Maunder, John E. 167–177
Medicinal plants 148–166
Meigs Co., Ohio 148–166
Melastomataceae 124–147
Meristem commitment and genet integrity 210–225
Mesoamerica 43–51
Miconia desportesii complex on Hispaniola 124–147; *M. desportesii* 129–133; *M. monciona* 134–136; *M. sphagnicola* 136–140; *M. tetrastoma* 140–145; key to species of *Miconia* 127–129; distribution maps 133, 139, 143
Mione, Thomas, Segundo Leiva G., and Leon Yacher. *Jaltomata andersonii* (Solanaceae): A new species of Peru. 118–123
Moore, Steven M. 197–209

Nautilocalyx biserrulatus, nom. nov. 46–47; ***N. purpurascens***, sp. nov. 43–45
NEBC Meeting News 85–89, 192—195, 296–301, 383
Neidium rudimentarum (Bacillariophyceae), a rare species with novel rudimentary canals and areolar openings from the temperate region of North America. 33–42

Neotropics, Bolivia 1–32
Neviusia dunthornei, sp. nov. 197–209
Neviusia leaves (Rosaceae: Kerrieae) from the lower-middle Eocene of southern British Columbia 197–209
New Books 84, 191, 295, 382
New England 97–117, 287–290
New Hampshire 60–65, 287–290, 359–366
New records: Maine 55–59, 287–290; Massachusetts 66–76; New England 287–290; New Hampshire 60–65, 287–290; Newfoundland 167–177
New species name: ***Nautilocalyx biserrulatus***, nom. nov. 46–47
New species: ***Jaltomata andersonii***, sp. nov. 118–123; ***Nautilocalyx purpurascens***, sp. nov. 43–45; ***Neviusia dunthornei***, sp. nov. 197–209; ***Paradrymonia alata***, sp. nov. 47–51
Newfoundland 167–177
North America, diatom 33–42
North Carolina 370–376
Notes on the box huckleberry, *Gaylussacia brachycera* (Ericaceae), and its unexpected presence in North Carolina. 370–376 (Note)
Noteworthy plants in created wetlands in southeastern New Hampshire. 60–65 (New England Note)

Ohio 148–166

Paradrymonia alata, sp. nov. 47–51
Patterns of plant invasions at sites with rare plant species throughout New England. 97–117
Penneys, Darin S. 124–147
Peru 118–123
Physalis species of southern U.S. 305–325; key to species 308–311; keys to varieties 314, 319
Phytodiversity, Bolivia 1–32
Pigg, Kathleen B. 197–209
Pinus strobus 273–286
Pityopsis, chromosome numbers 253–272
Plant communities of inland sandy acid pondshores in the lower Connecticut River Valley 66–76
Plant communities, Florida 226–252
Pringle, James S. Leptotypification of the name *Gentiana quinquefolia* var. *occidentalis* (Gentianaceae). 367–369 (Note)
Pseudosmodingium, taxonomic revision 347–358; *P. andrieuxii* 350–352; *P. barkleyi* 352–354; *P. perniciosum* 354–356; *P. virletii* 356–357; key to species 349; distribution map 351

Ramet 210–225
Rare plants: Massuchusetts 66–76; *Neidium rudimentarum* 33–42; New England 97–117; New Hampshire 60–65; Newfoundland 167–177
Records for *Carex rostrata* (Cyperaceae) in New England. 287–290 (New England Note)
Rediscovery of *Carex typhina* (Cyperaceae) in Maine. 52–54 (New England Note)
Reimer, Charles W. 33–42
Reviewers of Manuscripts 389
Revision of the species of the *Miconia desportesii* complex on Hispaniola 124–147
Rhizomes, *Dirca* spp. reproduction from 291–294
Ritter, Nur P. Too wet for aquatic plants? Floristic composition and phytodiversity in the wetlands along the base of the Bolivian Andes. 1–32
Rosaceae 197–209
Rudimentary canal, *Neidium rudimentarum* 33–42

Semple, John C. and Rachel E. Cook. Chromosome number determinations in fam. Compositae, tribe Astereae. VII. Mostly eastern North American and some Eurasian taxa. 253–272
Siver, Peter A. 33–42
Solanaceae 118–123; 305–325

Solidago, chromosome numbers 253–272
Sosa, Victoria 347–358
Southeastern United States 305–325
Sperduto, Dan 287–290
Statement of Ownership. No. 925, inside back cover
Sullivan, Janet R. The genus *Physalis* (Solanaceae) in the southeastern United States. 305–325
Symphyotrichum, chromosome numbers 253–272

Taxonomic revision of the genus *Pseudosmodingium* (Anacardiaceae). 347–358
Taxonomic studies in the Miconieae (Melastomataceae). VIII. A revision of the species of the *Miconia desportesii* complex on Hispaniola. 124–147
Thirty-six years of change in an Eastern Hemlock-White Pine stand in western Massachusetts. 273–286
Thymelaeaceae 291–294
Too wet for aquatic plants? Floristic composition and phytodiversity in the wetlands along the base of the Bolivian Andes. 1–32
Tsuga canadensis 273–286
Two new species and one new name in the Gesneriaceae from Costa Rica. 43–51

United Plant Savers (UpS), Ohio 148–166

Vascular flora, Ohio 148–166
Vascular plants of the Island of Newfoundland, Canada: Recent additions and rediscoveries. 167–177 (Note)
Vegetative growth versus flowering in *Maianthemum canadense* 210–225
Vegetative reproduction, *Dirca* spp. 291–294
Vernal pools on the Columbia Plateau of eastern Washington 326–346; species list 337–346

Washington 326–346
Wehr, Wesley C. 197–209
Wetlands, Bolivia 1–32; New Hampshire 60–65
Wilbur, Robert L. and Stefan Bloodworth. Notes on the box huckleberry, *Gaylussacia brachycera* (Ericaceae), and its unexpected presence in North Carolina. 370–376 (Note)

Yacher, Leon 118–123

Zebryk, Tad M. Inland sandy acid pondshores in the lower Connecticut River Valley, Hampden County, Massachusetts. 66–76 (New England Note)